

SIMPLE LIE COLOR ALGEBRAS OF WEYL TYPE¹

(To appear in *Israel J. Math.*)

Yucai Su², Kaiming Zhao and Linsheng Zhu²

Abstract. For an (ϵ, Γ) -color-commutative associative algebra A with an identity element over a field F of characteristic not 2, and for a color-commutative subalgebra D of color-derivations of A , denote by $A[D]$ the associative subalgebra of $\text{End}(A)$ generated by A (regarding as operators on A via left multiplication) and D . It is easily proved that, as an associative algebra, $A[D]$ is Γ -graded simple if and only if A is Γ -graded D -simple. Suppose A is Γ -graded D -simple. Then, (a) $A[D]$ is a free left A -module; (b) as a Lie color algebra, the subquotient $[A[D], A[D]]/Z(A[D]) \cap [A[D], A[D]]$ is simple (except one minor case), where $Z(A[D])$ is the color center of $A[D]$. The structure of this subquotient is explicitly described.

§1 Introduction

Lie color algebras are generalizations of Lie superalgebras. Let us start with the definition. Let F be a field, Γ an additive group. A *skew-symmetric bicharacter* of Γ is a map $\epsilon : \Gamma \times \Gamma \rightarrow F^*$ satisfying

$$\epsilon(\lambda, \mu) = \epsilon(\mu, \lambda)^{-1}, \quad \epsilon(\lambda, \mu + \nu) = \epsilon(\lambda, \mu)\epsilon(\lambda, \nu), \quad \forall \lambda, \mu, \nu \in \Gamma. \quad (1.1)$$

It is clear that $\epsilon(\lambda, 0) = 1$ for any $\lambda \in \Gamma$. Let $L = \bigoplus_{\lambda \in \Gamma} L_\lambda$ be a Γ -graded F -vector space. For a nonzero homogeneous element $a \in L$, denote by $\bar{a}$ the unique group element in Γ such that $a \in L_{\bar{a}}$. We shall call $\bar{a}$ the *color* of a . The F -bilinear map $[\cdot, \cdot] : L \times L \rightarrow L$ is called a Lie color bracket on L if the following conditions are satisfied:

$$[a, b] = -\epsilon(\bar{a}, \bar{b})[b, a], \quad (\text{skew symmetry}) \quad (1.2)$$

$$[a, [b, c]] = [[a, b], c] + \epsilon(\bar{a}, \bar{b})[b, [a, c]], \quad (\text{Jacoby identity}) \quad (1.3)$$

for all homogeneous elements $a, b, c \in L$. The algebra structure $(L, [\cdot, \cdot])$ is called an (ϵ, Γ) -Lie color algebra or simply a Lie color algebra. If $\Gamma = \mathbb{Z}/2\mathbb{Z}$ and $\epsilon(i, j) = (-1)^{ij}, \forall i, j \in \mathbb{Z}/2\mathbb{Z}$, then (ϵ, Γ) -Lie color algebras are simply Lie superalgebras. For Lie color algebras, we refer the reader to [P2] and the references there.

Let $A = \bigoplus_{\lambda \in \Gamma} A_\lambda$ be a Γ -graded associative F -algebra with an identity element 1, i.e., $A_\lambda A_\mu \subset A_{\lambda+\mu}$ for all $\lambda, \mu \in \Gamma$. So $1 \in A_0$. We say that A is *graded simple* if A does not have nontrivial Γ -graded ideals. Denote by $H(A)$ all homogeneous elements of A . If we define the bilinear product $[\cdot, \cdot]$ on A by

$$[x, y] = xy - \epsilon(\bar{x}, \bar{y})yx, \quad \forall x, y \in H(A), \quad (1.4)$$

then $(A, [\cdot, \cdot])$ becomes a (ϵ, Γ) -Lie color algebra. We shall simply write $\epsilon(x, y)$ for $\epsilon(\bar{x}, \bar{y})$ for $x, y \in H(A)$.

¹AMS Subject Classification - Primary: 17B20, 17B65, 17B67, 17B68, 17B75, 16A68, 16A40.

This work was supported by NSF of China, National Educational Department of China, Jiangsu Educational Committee, and Hundred Talents Program of Chinese Academy of Sciences.

²These authors were partially supported by Academy of Mathematics and System Sciences during their visit to this academy.

An *Lie color ideal* U of A is a Γ -graded vector space U of A such that $[A, U] \subset U$. Sometimes it is called an (ϵ, Γ) -Lie ideal, or simply a color ideal. The ϵ -center $Z(A)$ of A is defined as

$$Z = Z(A) = \{x \in A \mid [x, A] = 0\}. \quad (1.5)$$

It is easy to see that $Z(A)$ is Γ -graded. We say that A is *color-commutative* (or ϵ -color commutative) if $[A, A] = 0$.

Passman [P1, P2] proved that, for a color-commutative associative algebra A with an identity element over a field F , and for a color-commutative color-derivation subalgebra D of A , the Lie color algebra (including the Lie algebra case) $AD = A \otimes D$ is simple if and only if A is graded D -simple and AD acts faithfully on A and, $\text{char } F \neq 2$ or $\dim_F D \geq 2$ or $D(A) = A$. In [SZ1, SZ2, Z], (associative and Lie) algebras of Weyl type were constructed and studied. In this paper we shall study the color version of these algebras.

For a color-commutative associative algebra A with an identity element over a field F of characteristic not 2, and for a color-commutative subalgebra D of color-derivations of A , denote by $A[D]$ the associative subalgebra of $\text{End}(A)$ generated by A (regarding as operators on A via multiplication) and D . It is easily proved that, as an associative algebra, $A[D]$ is graded simple if and only if A is graded D -simple (see Theorem 2.3). Suppose A is graded D -simple. Then, (a) $A[D]$ is a free left A -module; (b) as a Lie color algebra, the subquotient $[A[D], A[D]]/Z(A[D]) \cap [A[D], A[D]]$ is simple (except one minor case), where $Z(A[D])$ is the ϵ -center of $A[D]$. The structure of this subquotient is explicitly described (see Theorem 3.9). In many cases $A[D] = [A[D], A[D]]$.

§2 Graded simple associative algebras of Weyl type

Throughout this work, we assume that A is Γ -graded and (ϵ, Γ) -color commutative and the field F is of characteristic not 2. An F -linear transformation $\partial : A \rightarrow A$ is called a *homogeneous color-derivation* of degree $\bar{\partial} \in \Gamma$ if

$$\begin{aligned} \partial(a) &\in A_{\bar{\partial} + \bar{a}}, \quad \forall a \in H(A) \text{ and} \\ \partial(ab) &= \partial(a)b + \epsilon(\partial, a)a\partial(b), \quad \forall a, b \in H(A). \end{aligned} \quad (2.1)$$

Taking $a = b = 1$ in (2.1), we obtain $\partial(1) = 0$ and so $\partial(c) = 0$ for all $c \in F$. Denote $\text{Der}^\epsilon(A) = \bigoplus_{\lambda \in \Gamma} \text{Der}_\lambda^\epsilon(A)$, where $\text{Der}_\lambda^\epsilon(A)$ is the F -vector space spanned by all homogeneous color derivations of degree $\lambda \in \Gamma$. As in the Lie algebras case, it is easy to verify that $\text{Der}^\epsilon(A)$ becomes an (ϵ, Γ) -Lie color algebra under the Lie color bracket

$$[\partial, \partial'] = \partial\partial' - \epsilon(\partial, \partial')\partial'\partial, \quad \forall \partial, \partial' \in H(\text{Der}^\epsilon(A)), \quad (2.2)$$

where $\partial\partial'$ is the composition of the operators ∂ and ∂' .

Let $D = \bigoplus_{\lambda \in \Gamma} D_\lambda$ be a nonzero Γ -graded color-commutative Lie color subalgebra of $\text{Der}^\epsilon(A)$, i.e.,

$$\partial\partial' = \epsilon(\partial, \partial')\partial'\partial, \quad \forall \partial, \partial' \in H(D). \quad (2.3)$$

We call A is *graded D -simple* if A has no nontrivial graded D -stable ideals (see [P2]). Set $\Gamma_+ = \{\lambda \in \Gamma \mid \epsilon(\lambda, \lambda) = 1\}$. Then by (1.1), Γ_+ is a subgroup of Γ with index ≤ 2 . Set

$\Gamma_- = \{\lambda \in \Gamma \mid \epsilon(\lambda, \lambda) = -1\}$. For any graded subspace B of A , we define $B_+ = \bigoplus_{\lambda \in \Gamma_+} B_\lambda$, then B_+ is Γ -graded. Similarly we can define B_- . Since $\Gamma = \Gamma_+ \cup \Gamma_-$, it follows that $B = B_+ \oplus B_-$. By (2.3), we have $\partial^2 = 0$ if $\partial \in H(D_-)$.

Fix a homogeneous basis $\{\partial_i \mid i \in I\}$ of D , where I is some index set. Fix a total ordering $<$ on I . Define J to be the set of all $\alpha = (\alpha_i \mid i \in I) \in \mathbb{Z}_+^I$ such that $\alpha_i = 0$ for all but a finite number of $i \in I$ and $\alpha_i = 0$ or 1 if $\bar{\partial}_i \in \Gamma_-$. For $\alpha \in J$, we define the *support* of α by $\text{supp}(\alpha) = \{i \in I \mid \alpha_i \neq 0\}$ and define the *level* of α by $l(\alpha) = |\alpha| = \sum_{i \in I} \alpha_i$. Define a total ordering on J by

$$\alpha < \beta \Leftrightarrow |\alpha| < |\beta|, \text{ or } |\alpha| = |\beta| \text{ and } \exists i \in I \text{ such that } \alpha_i < \beta_i \text{ and } \alpha_j = \beta_j, \forall j \in I, j < i. \quad (2.4)$$

Denote by $F[D]$ the Γ -graded associative color-commutative algebra generated by D , and denote by $A[D]$ the associative subalgebra of $\text{End}(A)$ generated by A (regarding as operators on A via left multiplication) and D . For $x \in A, \alpha \in J$, we define by $x\partial^\alpha$ the operator acting on A , i.e.,

$$(x\partial^\alpha)(y) = x(\partial_1^{\alpha_1}(\partial_2^{\alpha_2} \dots (\partial_r^{\alpha_r}(y)) \dots)), \forall y \in A, \quad (2.5)$$

where $\partial^\alpha = \partial_{i_1}^{\alpha_{i_1}} \partial_{i_2}^{\alpha_{i_2}} \dots \partial_{i_r}^{\alpha_{i_r}}$ with $i_1 < i_2 < \dots < i_r$. We shall call α the *degree* of ∂^α . Define $\epsilon^+(\alpha, \beta)$ for $\alpha, \beta \in \Gamma$ by $\partial^\alpha \partial^\beta = \epsilon^+(\alpha, \beta) \partial^{\alpha+\beta}$, then

$$\epsilon^+(\alpha, \beta) = \prod_{i, j \in I: i > j} \epsilon(\partial_i, \partial_j)^{\alpha_i \beta_j}, \forall \alpha, \beta \in J. \quad (2.6)$$

We shall also simply write $\epsilon^+(x, y)$ for $\epsilon^+(\bar{x}, \bar{y})$ for $x, y \in H(A[D])$. From $\epsilon^+(\alpha, \beta) \partial^{\alpha+\beta} = \partial^\alpha \partial^\beta = \epsilon(\alpha, \beta) \partial^\beta \partial^\alpha = \epsilon(\alpha, \beta) \epsilon^+(\beta, \alpha) \partial^{\alpha+\beta}$, we see that

$$\epsilon(\alpha, \beta) = \epsilon^+(\alpha, \beta) \epsilon^+(\beta, \alpha)^{-1}, \forall \alpha, \beta \in J, \quad (2.7)$$

If $\alpha + \beta \notin J$, we set $\partial^{\alpha+\beta} = 0$. For any $\alpha \in J$, set

$$J(\alpha) = \{\gamma \in J \mid \gamma_i \leq \alpha_i, \forall i \in I\}. \quad (2.8)$$

Denote $\binom{\alpha}{\gamma} = \prod_{i \in I} \binom{\alpha_i}{\gamma_i}$, $\forall \alpha, \gamma \in J, x \in H(A)$. Then $\binom{\alpha}{\gamma} = 0$ if $\gamma \notin J(\alpha)$. We have

$$\partial^\alpha(xy) = \sum_{\gamma \in J} \binom{\alpha}{\gamma} \epsilon^+(\alpha - \gamma, \gamma)^{-1} \epsilon(\partial^\gamma, x) \partial^{\alpha-\gamma}(x) \partial^\gamma(y), \quad \forall \alpha \in J, x, y \in H(A). \quad (2.9)$$

One may easily check that

$$(u\partial^\alpha) \cdot (v\partial^\beta) = \sum_{\lambda \in J} \binom{\alpha}{\lambda} \epsilon^+(\alpha - \lambda, \lambda)^{-1} \epsilon(\partial^\lambda, v) \epsilon^+(\lambda, \beta) u \partial^{\alpha-\lambda}(v) \partial^{\beta+\lambda}, \quad \forall u, v \in H(A), \alpha, \beta \in J. \quad (2.10)$$

Clearly that formula (2.10) defines a Γ -graded associative algebra $(A[D], \cdot)$ such that A is a left $A[D]$ -module via (2.5). For any

$$x = \sum_{\alpha \in J} u_\alpha \partial^\alpha \in A[D], \quad u_\alpha \in A, \quad (2.11)$$

the expression in (2.11) is sometimes not unique. An expression of x in (2.11) is called *principal*, if the integer $\max\{|\alpha| \mid a_\alpha \neq 0\}$ is minimal. We denote this integer by $h(x)$. Set

$$F_1 = A^D = \{u \in A \mid D(u) = 0\}. \quad (2.12)$$

From Lemma 2.1 in [P2], any nonzero $a \in H(F_1)$ is invertible (i.e., F_1 is a graded field) when A is Γ -graded D -simple. In this case $(F_1)_- = 0$.

Lemma 2.1. *i) $A \cap A[D]D = 0$. ii) The ϵ -center $Z(A[D])$ of $A[D]$ is F_1 .*

Proof. Note that we have assume that A is Γ -graded (ϵ, Γ) -commutative. Part i) follows from the action of $x \in A \cap A[D]D$ on 1.

ii). For all $u \in H(F_1)$ and $\delta \in H(D)$, we have $[\delta, u] = \delta(u) = 0$. So $F_1 \subset Z(A[D])$. Suppose $x \in H(Z(A[D]))$. If $h(x) > 0$, write $x = x_0 + x_1$ where $x_0 \in H(A)$ and $x_1 \in H(A[D]D)$. Then $x_1 \neq 0$. Choose $a \in H(A)$ such that $x_1(a) \neq 0$. It follows that $0 = [x, a] = [x_1, a] = x_1(a) + y$, where $y \in A[D]D$. From i) we deduce that $x_1(a) = y = 0$, a contradiction. So $h(x) = 0$, i.e., $x \in A$. We have

$$x\partial = \epsilon(\bar{x}, \bar{\partial})\partial x = \epsilon(\bar{x}, \bar{\partial})\partial(x) + x\partial, \quad \forall \partial \in H(D),$$

to give $\partial(x) = 0$ for all $\partial \in H(D)$, i.e., $x \in F_1$. Therefore Lemma 2.1 follows. $\square$

Theorem 2.2. *Suppose that A is a Γ -graded (ϵ, Γ) -commutative F -algebra with an identity, and $D \subset \text{Der}^\epsilon(A)$ is a color commutative subspace. Then the Γ -graded associative algebra $A[D]$ is graded simple if and only if A is Γ -graded D -simple.*

Proof. " $\Rightarrow$ ": Suppose A is not graded D -simple. Choose a nonzero proper graded D -stable ideal $\mathcal{K}$. Then clearly $\mathcal{K}[D]$ is a nonzero graded color ideal of $A[D]$. Since $A[D]$ is graded simple, then $\mathcal{K}[D] = A[D]$, in particular, $A \subset \mathcal{K}[D]$. From lemma 2.1 i), we know that $A \subset \mathcal{K}$, a contradiction. Thus A is graded D -simple.

" $\Leftarrow$ ": Suppose L is a nonzero graded ideal of $A[D]$. It suffices to show that $L = A[D]$.

Suppose that $L \cap A = 0$. Choose $x \in H(L) \setminus \{0\}$ such that $h(x)$ is minimal. So $h(x) > 0$. Write $x = x_0 + x_1$ where $x_0 \in H(A)$ and $x_1 \in H(A[D]D)$. Then $x_1 \neq 0$. Choose $a \in H(A)$ such that $x_1(a) \neq 0$. It follows that

$$x' = [x, a] = [x_1, a] = x_1(a) + y \in L,$$

where $y \in H(A[D]D)$. From (1.4) and the computation (2.10), we deduce that $h(x') = h(y) < h(x)$. By the minimality of $h(x)$ we deduce that $x' = 0$. Applying Lemma 2.2 i) gives $x_1(a) = y = 0$, a contradiction. Thus $L \cap A \neq 0$. It is clear that $L \cap A$ is a graded D -ideal of A . Since A is graded D -simple, $L \cap A = A$. In particular $1 \in L$. Therefore $L = A[D]$. $\square$

§3 Simple Lie color algebras of Weyl type

In this section we assume that A is Γ -graded D -simple and (ϵ, Γ) -commutative associative algebra with an identity, the field F is of characteristic not 2, and $D \subset \text{Der}^\epsilon(A)$ is color commutative. In this section we shall first study the structure of $A[D]$ as a left

A -module, then investigate the Lie structure of $A[D]$. We still use the notation in Section 2.

Lemma 3.1. *i) If $xd^\alpha = 0$ for some $x \in H(A) \setminus \{0\}$ and some $\alpha \in J$, then $d^\alpha = 0$.
ii) $A \subset [A[D], A[D]]$.*

Proof. i) Let $C = \{x \in A \mid x\partial^\alpha = 0\}$. It is easy to verify that C is a nonzero Γ -graded D -ideal of A . Since A is Γ -graded D -simple, then $C = A$. It yields that $\partial^\alpha = 0$.

ii) From $[x\partial, y] = x\partial(y) \in [A[D], A[D]]$ for all $x, y \in H(A)$ and $\partial \in H(D)$, we obtain that $AD(A) \subset [A[D], A[D]]$. Since $AD(A)$ is a nonzero Γ -graded D -ideal of A and A is Γ -graded D -simple, $A = AD(A)$. Thus $A \subset [A[D], A[D]]$. $\square$

If $\text{char } F = p > 0$, observe that for any $\partial \in \text{Der}^\epsilon(A)$, one has $\partial^p \in \text{Der}^\epsilon(A)$. For convenience, no matter whether $\text{char } F = p > 0$ or $\text{char } F = 0$, we denote

$$\mathcal{D} = \text{Der}^\epsilon(A) \cap F_1[D], \quad (3.1)$$

where $F_1[D]$ is the subalgebra of $\text{End}(A)$ generated by F_1 and D . Let $\{d_i \mid i \in \mathcal{I}\}$ be a homogeneous F_1 -basis for $\mathcal{D}$, where $\mathcal{I}$ is some index set. It is clear that $D \subset \mathcal{D}$, and that A is graded D -simple if and only if A is graded $\mathcal{D}$ -simple. Let

$$\begin{aligned} \mathcal{J} = \{ \alpha = (\alpha_i \mid i \in \mathcal{I}) \mid \alpha_i \in \mathbb{Z}_+, \text{ and } \alpha_i \leq p-1 \text{ if } \text{char } F = p > 0, \\ \text{and } \alpha_i = 0 \text{ or } 1 \text{ if } \bar{d}_i \in \Gamma_-, \text{ and } \alpha_i = 0 \text{ for all but a finite number of } i \in \mathcal{I} \}. \end{aligned} \quad (3.2)$$

We will often simply denote $\alpha = (\alpha_i \mid i \in \mathcal{I})$ by $\alpha = (\alpha_i)$. We also fix a total ordering $<$ on $\mathcal{I}$ and define a total ordering on $\mathcal{J}$ as in (2.7). We also write $d^\alpha = \prod_{i \in \mathcal{I}} d^{\alpha_i}$ according to the ordering $<$ on $\mathcal{I}$ for $\alpha \in \mathcal{J}$. Then

$$A[D] = \sum_{\alpha \in \mathcal{J}} Ad^\alpha. \quad (3.3)$$

Theorem 3.2. *Suppose that the field F is of characteristic not 2, A is a Γ -graded D -simple and (ϵ, Γ) -commutative associative F -algebra with an identity, where $D \subset \text{Der}^\epsilon(A)$ is a color commutative subspace. Then $A[D]$ is a free A -module with the homogeneous basis $\{d^\alpha \mid \alpha \in \mathcal{J}\}$.*

We break the proof of this theorem into several Lemmas. For $\mathcal{J}_0 \subset \mathcal{J}$, $B \subset A$, we say that the sum $\sum_{\alpha \in \mathcal{J}_0} Bd^\alpha$ is *direct* if a finite sum $\sum_{\alpha \in \mathcal{J}_0} b_\alpha d^\alpha = 0$ with $b_\alpha \in B$ implies $b_\alpha d^\alpha = 0$ for all $\alpha \in \mathcal{J}_0$.

Lemma 3.3. *Let $\mathcal{J}_0 \subset \mathcal{J}$. The sum $\sum_{\alpha \in \mathcal{J}_0} Ad^\alpha$ is direct if and only if the sum $\sum_{\alpha \in \mathcal{J}_0} F_1 d^\alpha$ is direct.*

Proof. “ $\Rightarrow$ ”: This direction is clear.

“ $\Leftarrow$ ”: Suppose $\sum_{\alpha \in \mathcal{J}_0} Ad^\alpha$ is not direct. There exists $\mathcal{J}_1 = \{\alpha^{(0)}, \alpha^{(1)}, \dots, \alpha^{(r)}\} \subset \mathcal{J}_0$ with $|\mathcal{J}_1| > 1$ such that $Ad^{\alpha^{(0)}} \cap \sum_{i=1}^r Ad^{\alpha^{(i)}} \neq 0$. Choose such a minimal r . Let

$$ud^{\alpha^{(0)}} = \sum_{i=1}^r u_i d^{\alpha^{(i)}} \in (Ad^{\alpha^{(0)}} \cap \sum_{i=1}^r Ad^{\alpha^{(i)}}) \setminus \{0\}, \quad (3.4)$$

for some $u, u_i \in H(A)$, and let $I_1 = \{0, 1, \dots, r\}$. It follows that $d^{\alpha^{(i)}} \neq 0$ for any $i \in I_1$. Let

$$B_0 = \{x \in A \mid x d^{\alpha^{(0)}} \in \sum_{i \in I_1 \setminus \{0\}} A d^{\alpha^{(i)}}\}.$$

Then it is easy to see that B_0 is a nonzero Γ -graded D -ideal of A . Since A is Γ -graded D -simple, then $B_0 = A$. Then

$$d^{\alpha^{(0)}} = \sum_{i \in I_1 \setminus \{0\}} a_i d^{\alpha^{(i)}} \text{ for some } a_i \in A. \quad (3.5)$$

By taking bracket with $\partial \in H(\mathcal{D})$, we deduce that

$$0 = [\partial, d^{\alpha^{(0)}}] = [\partial, \sum_{i \in I_1 \setminus \{0\}} a_i d^{\alpha^{(i)}}] = \sum_{i \in I_1 \setminus \{0\}} \partial(a_i) d^{\alpha^{(i)}}.$$

From the minimality of r , we must have

$$\partial(a_i) d^{\alpha^{(i)}} = 0, \forall \partial \in \mathcal{D}, 1 \leq i \leq r. \quad (3.6)$$

Suppose $D(a_i) \neq 0$ for some $i \in I_1$, say $D(a_1) \neq 0$. Lemma 3.1 yields that $d^{\alpha^{(1)}} = 0$, a contradiction. Thus $D(a_i) = 0$ for all $i \in I_1$, i.e., $a_i \in F_1$ in (3.5). Therefore the sum $\sum_{\alpha \in \mathcal{J}_1} F_1 d^\alpha$ is not direct. This proves the lemma. $\square$

Lemma 3.4. $A[D] = \oplus_{\alpha \in \mathcal{J}} A d^\alpha$.

Proof. Suppose the sum in (3.3) is not direct. From Lemmas 2.1 i) and 3.1, there exists $\mathcal{J}_1 = \{\alpha^{(0)}, \alpha^{(1)}, \dots, \alpha^{(r)}\} \subset \mathcal{J} \setminus \{0\}$ such that $F_1 d^{\alpha^{(0)}} \cap \sum_{i=1}^r F_1 d^{\alpha^{(i)}} \neq 0$. Let $I_1 = \{0, 1, \dots, r\}$, and let $h(\mathcal{J}_1) = \max\{|\alpha^{(i)}| \mid i \in I_1\}$. Then $h(\mathcal{J}_1) > 1$. We choose $\mathcal{J}_1$ such that $h(\mathcal{J}_1)$ is minimal, and then r is minimal. Let

$$-d^{\alpha^{(0)}} = \sum_{i=1}^r a_i d^{\alpha^{(i)}} \in (F_1 d^{\alpha^{(0)}} \cap \sum_{i=1}^r F_1 d^{\alpha^{(i)}}) \setminus \{0\}, \quad (3.7)$$

for some $a_i \in H(F_1)$, $1 \leq i \leq r$. Denote $a_0 = 1$. It follows that $a_i d^{\alpha^{(i)}} \neq 0$ for $i \in I_1$. Rewrite (3.7) to give $\sum_{i=0}^r a_i d^{\alpha^{(i)}} = 0$. Then for $x \in H(A)$, $\sum_{i=0}^r a_i d^{\alpha^{(i)}}(x) = 0$. By (2.10),

$$\begin{aligned} 0 &= [\sum_{i=0}^r a_i d^{\alpha^{(i)}}, x] = \sum_{i=0}^r \sum_{\gamma \in J} a_i (\alpha_\gamma^{(i)}) \epsilon^+(\alpha^{(i)} - \gamma, \gamma)^{-1} \epsilon(d^\gamma, x) d^{\alpha^{(i)} - \gamma}(x) d^\gamma \\ &= \sum_{\gamma \in J, 0 < |\gamma| < h(\mathcal{J}_1)} \left(\sum_{i=0}^r a_i (\alpha_\gamma^{(i)}) \epsilon^+(\alpha^{(i)} - \gamma, \gamma)^{-1} \epsilon(d^\gamma, x) d^{\alpha^{(i)} - \gamma}(x) \right) d^\gamma. \end{aligned}$$

From the minimality of $h(\mathcal{J}_1)$ and Lemma 2.1 i), it follows that

$$\sum_{i=0}^r a_i (\alpha_\gamma^{(i)}) \epsilon^+(\alpha^{(i)} - \gamma, \gamma)^{-1} \epsilon(d^\gamma, x) d^{\alpha^{(i)} - \gamma}(x) d^\gamma = 0, \quad (3.8)$$

for all $x \in H(A)$ and all $\gamma \in \mathcal{J} \setminus \{0\}$ with $0 < |\gamma| < h(\mathcal{J}_1)$. Noting that for any $\gamma \in \mathcal{J}$ with $\alpha^{(i)} - \gamma \in \mathcal{J}$, we have $d^\gamma d^{\alpha^{(i)} - \gamma} = \epsilon(d^\gamma, d^{\alpha^{(i)} - \gamma}) d^{\alpha^{(i)}} \neq 0$, to give $d^\gamma \neq 0$. Since $d^\gamma \neq 0$, from (3.8) and Lemma 3.1 we obtain $\sum_{i=0}^r a_i(\alpha_\gamma^{(i)}) \epsilon^+(\alpha^{(i)} - \gamma, \gamma)^{-1} \epsilon(d^\gamma, x) d^{\alpha^{(i)} - \gamma}(x) = 0$, i.e.,

$$\sum_{i=0}^r a_i(\alpha_\gamma^{(i)}) \epsilon^+(\alpha^{(i)} - \gamma, \gamma)^{-1} d^{\alpha^{(i)} - \gamma}(\epsilon(d^\gamma, x)x) = 0,$$

for all $x \in H(A)$ and all $\gamma \in \mathcal{J}$ with $\alpha^{(i)} - \gamma \in \mathcal{J}$. So $\sum_{i=0}^r a_i(\alpha_\gamma^{(i)}) \epsilon^+(\alpha^{(i)} - \gamma, \gamma)^{-1} d^{\alpha^{(i)} - \gamma} = 0$ for all $\gamma \in \mathcal{J}$ with $0 < h(\gamma) < h(\mathcal{J}_1)$. We have $a_i(\alpha_\gamma^{(i)}) \neq 0$ since $\alpha_j^{(i)} \leq p - 1$ if $\text{char } F = p > 0$, for $\gamma \in \mathcal{J}$ with $\alpha^{(i)} - \gamma \in \mathcal{J}$. Thus we see that $d^{\alpha^{(i)} - \gamma} = 0$ for all $\gamma \in \mathcal{J} \setminus \{0\}$ with $\alpha^{(i)} - \gamma \in \mathcal{J} \setminus \{0\}$. As we noticed also $d^{\alpha^{(i)} - \gamma} \neq 0$, so this contradicts the minimality of $h(\mathcal{J}_1)$. Therefore $A[D] = \bigoplus_{\alpha \in \mathcal{J}_0} A d^\alpha$. $\square$

Lemma 3.5. *For all $\alpha \in \mathcal{J}$, $d^\alpha \neq 0$.*

Proof. It is obvious that $d^\alpha \neq 0$ for $\alpha \in \mathcal{J}$ with $|\alpha| \leq 1$. Suppose $d^\alpha = 0$ for some $\alpha \in \mathcal{J}$ with $|\alpha| \geq 2$. Choose α such that $|\alpha|$ is minimal. Then for any $x \in H(A)$, $d^\alpha(x) = 0$. Then by (1.4), (2.10),

$$0 = [d^\alpha, x] = \sum_{\beta \in \mathcal{J}} \binom{\alpha}{\beta} \epsilon^+(\alpha - \beta, \beta)^{-1} \epsilon(\beta, \bar{x}) d^{\alpha - \beta}(x) d^\beta, \forall x \in H(A).$$

Applying Lemma 3.4 and noting that $\binom{\alpha}{\beta} \epsilon^+(\alpha - \beta, \beta)^{-1} \epsilon(\beta, \bar{x}) \neq 0$ (by noting that $\alpha_i \leq p - 1$ if $\text{char } F = p > 0$) for any $\beta \in \mathcal{J}$ with $\alpha - \beta \in \mathcal{J}$, we get $d^{\alpha - \beta}(x) d^\beta = 0, \forall x \in H(A), \beta \in \mathcal{J}$. Noting that $d^\beta \neq 0$ for any $\beta \in \mathcal{J}$ with $\alpha - \gamma \in \mathcal{J} \setminus \{0\}$ (since $|\alpha|$ is minimal), applying Lemma 3.1 gives $d^{\alpha - \beta}(x) = 0, \forall x \in H(A)$, i.e., $d^{\alpha - \beta} = 0$ for $\beta \in \mathcal{J}$ with $\alpha - \gamma \in \mathcal{J} \setminus \{0\}$, contradicting the minimality of $|\alpha|$. Therefore the assertion follows. $\square$

Proof of Theorem 3.2. It follows from Lemmas 3.1, 3.3, 3.4 and 3.5. $\square$

Denote by $W = [A[D], A[D]]$ the derived Lie color ideal of $A[D]$. In the rest of this section We will investigate the structure of W .

Lemma 3.6. *Suppose $\dim_{F_1} \mathcal{D} > 1$ or $\mathcal{D} \neq \mathcal{D}_-$. Then for any homogeneous color-derivation $\partial \in \mathcal{D}$, we have $A\partial \subset W$.*

Proof. Case 1: $\partial \in \mathcal{D}_+$. For any $x, a \in H(A)$, by Lemma 3.1 ii), $a, x\partial^2(a) \in W$, thus

$$x\epsilon(\partial, a)\partial(a)\partial = \frac{1}{2}([x\partial^2, a] - x\partial^2(a)) \in W, \quad (3.9)$$

i.e., $A\partial(A)\partial \subset W$. But the space $K = \{x \in A \mid x\partial \in W\}$ is Γ -graded D -stable since W is a Lie color ideal of $A[D]$, and K contains the nonzero graded ideal $A\partial(A)$ of A , by Lemma 3.1 in [P2], $K = A$, i.e., $A\partial \subset W$.

Case 2: $\partial \in \mathcal{D}_-$. First assume that $\mathcal{D}_+ \neq 0$. Choose $\partial' \in H(\mathcal{D}_+) \setminus \{0\}$, by Case 1, $A\partial' \subset W$. Thus for any $x, y \in H(A)$, we have $[x\partial', y\partial] \in W$ and

$$[x\partial', y\partial] = x\partial'(y)\partial - \epsilon(x\partial', y\partial)y\partial(x)\partial' \equiv x\partial'(y)\partial \pmod{W}, \quad (3.10)$$

i.e., $A\partial'(A)\partial \subset W$, thus as in Case 1, $A\partial \subset W$. Next assume that $\mathcal{D}_+ = 0$, then by the assumption of the lemma, we can choose $\partial' \in \mathcal{D}_-$ such that ∂, ∂' are F_1 -linear independent. For $x, y \in H(A)$, we have $[x\partial', y\partial] \in W$ and

$$\begin{aligned} [x\partial', y\partial] &= x\partial'(y)\partial - \epsilon(x\partial', y\partial)y\partial(x)\partial' \\ &= x\partial'(y)\partial - \epsilon(x\partial', y\partial)\epsilon(\partial, y)^{-1}(\partial(yx)\partial' - \partial(y)x\partial') \\ &\equiv x(\partial'(y)\partial + \epsilon(x\partial', y\partial)\epsilon(y, \partial)\epsilon(\partial(y), x)\partial(y)\partial') \pmod{W} \\ &= x(\partial'(y)\partial + \epsilon(\partial', y\partial)\epsilon(y, \partial)\partial(y)\partial'), \end{aligned} \quad (3.11)$$

where the equality “ $\equiv$ ” follows from that $\partial(xy)\partial' = [\partial, xy\partial'] \in W$. Denote $d = \partial'(y)\partial + \epsilon(\partial', y\partial)\epsilon(y, \partial)\partial(y)\partial'$. Note that (3.11) shows that $Ad \in W$. By using $\partial^2 = 0$ and applying $\text{ad } \partial$ to (3.11), we obtain $[\partial, xd] \in W$ and

$$[\partial, xd] = \partial(x)d + \epsilon(\partial, x)x[\partial, d] \equiv \epsilon(\partial, x)x\partial(\partial'(y))\partial \pmod{W}.$$

This shows that $A\partial(\partial'(A))\partial \subset W$. But $\partial\partial' \neq 0$ by Lemma 3.5, thus as in Case 1, $A\partial \subset W$. $\square$

For any $i \in \mathcal{I}$, we define

$$\delta^{(i)} \in \mathcal{J} \text{ such that } \delta_j^{(i)} = \delta_{i,j}, \forall j \in \mathcal{I}. \quad (3.12)$$

The following technical lemma plays a crucial role in describing the structure of W .

Lemma 3.7. *Let $\beta \in \mathcal{J}$ with $\text{supp}(\beta) = \{1, 2, \dots, n\}$ and let $\partial'_1, \partial'_2, \dots, \partial'_n \in \text{Der}^\epsilon(A)$ be A -linear independent homogeneous derivations such that $\bar{\partial}'_i - \bar{d}_i = \bar{\partial}'_j - \bar{d}_j$ for $i, j = 1, 2, \dots, n$. Suppose there exist $a_1, a_2, \dots, a_n \in F \setminus \{0\}$ such that*

$$x \sum_{i=1}^n a_i \epsilon(d^{\beta-\delta^{(i)}}) \partial'_i(b) d^{\beta-\delta^{(i)}} \in W, \forall x, b \in H(A), \quad (3.13)$$

then $Ad^{\beta-\delta^{(i)}} \subset W$ for $i = 1, 2, \dots, n$.

Proof. By shifting the index, it suffices to prove $Ad^{\beta-\delta^{(n)}} \subset W$. We shall prove by induction on n . If $n = 1$, (3.13) shows that $A\partial'_1(A)d^{\beta-\delta^{(n)}} \subset W$, thus the result follows from [P2]. Suppose $n \geq 2$. Replacing x by $x\partial'_1(a)$, and replacing x, b by $y\partial'_1(b), a$ in (3.13), we obtain respectively

$$\begin{aligned} x \sum_{i=1}^n a_i \epsilon(d^{\beta-\delta^{(i)}}) \partial'_1(a) \partial'_i(b) d^{\beta-\delta^{(i)}} &\in W, \\ y \sum_{i=1}^n a_i \epsilon(d^{\beta-\delta^{(i)}}) \partial'_1(b) \partial'_i(a) \partial'_i(b) d^{\beta-\delta^{(i)}} &\in W. \end{aligned} \quad (3.14)$$

Setting $y = x\epsilon(d^{\beta-\delta^{(1)}}) \partial'_1(a)^{-1} \epsilon(d^{\beta-\delta^{(1)}}) \partial'_1(b) \partial'_1(a)^{-1}$ and subtracting the two expressions of (3.14), we obtain

$$x \sum_{i=2}^n a_i \epsilon(d^{\beta-\delta^{(i)}}) (\partial'_1(a) \partial'_i(b) - \partial'_i(a) \partial'_1(b)) d^{\beta-\delta^{(i)}} \in W, \quad (3.15)$$

where

$$\begin{aligned}
u_i(a) &= \epsilon(d^{\beta-\delta^{(1)}}, a)^{-1} \epsilon(d^{\beta-\delta^{(1)}}, b) \epsilon(\partial'_1(b), \partial'_1(a))^{-1} \\
&\quad \cdot \epsilon(d^{\beta-\delta^{(i)}}, a)^{-1} \epsilon(d^{\beta-\delta^{(i)}}, b) \epsilon(\partial'_1(b), \partial'_i(a)) \\
&= \epsilon(d^{\delta^{(1)}-\delta^{(i)}}, a) \epsilon(\partial'_1, \partial'_i \partial'_1) \in F.
\end{aligned} \tag{3.16}$$

Fix $a \in H(A)$ such that $\partial'_1(a) \neq 0$, then $\partial''_i = \partial'_1(a) \partial'_i - u_i(a) \partial'_i(a) \partial'_1 \in \text{Der}_F^\epsilon(A)$, $i = 2, \dots, n$ are A -linear independent derivations satisfying the conditions of the Lemma, thus by induction $Ad^{\beta-\delta^{(n)}} \subset W$. $\square$

Lemma 3.8 [ZS, Theorem 2.12]. *Suppose that A is a Γ -graded simple associative algebra of characteristic not 2 and $\dim_{Z(A)} A > 4$. Then $[A, A]/([A, A] \cap Z)$ is a simple ϵ -Lie color algebra.* $\square$

Our second main result in this section is the following.

Theorem 3.9. *Suppose that F is of characteristic not 2, A is a Γ -graded D -simple (ϵ, Γ) -commutative associative F -algebra, where $D \subset \text{Der}^\epsilon(A)$ is color commutative and nonzero. Let $W = [A[D], A[D]]$.*

i) *If $|\mathcal{J}| = \infty$, then $W = A[D]$.*

ii) *Suppose $|\mathcal{J}| < \infty$. Let $\gamma \in \mathcal{J}$ be the maximal element of $\mathcal{J}$ (i.e., $|\gamma| = h(\mathcal{J})$), then $W = \oplus_{\alpha \in \mathcal{J} \setminus \{\gamma\}} Ad^\alpha \oplus \mathcal{D}(A)d^\gamma$.*

iii) *The Lie color algebra $\overline{A[D]} = W/F_1$ is simple except when $A = F_1[t]/(t^2)$ and $D = F \frac{d}{dt}$ and both t and $\frac{d}{dt}$ have colors in Γ_- .*

Proof. If $\dim \mathcal{D} = 1$ and $\mathcal{D} = \mathcal{D}_-$, we see that $|\mathcal{J}| = 2$. Clearly ii) is true in this case. Now suppose that $\mathcal{D} \neq \mathcal{D}_-$ if $\dim \mathcal{D} = 1$. Then $|\mathcal{J}| > 2$.

For any $\beta \in \mathcal{J}$ with $|\beta| \geq 2$. Using inductive assumption, we may suppose that $Ad^\alpha \subset W$ for all $\alpha \in \mathcal{J}(\beta)$ with $|\alpha| \leq |\beta| - 2$. Assume that $\text{supp}(\beta) = \{1, 2, \dots, n\}$. Then for all $x, b \in H(A)$ we have $[xd^\beta, b] \in W$ and

$$[xd^\beta, b] = x[d^\beta, a] \equiv x \sum_{i=1}^n \beta_i \epsilon(d^{\beta-\delta^{(1)}}, b) \epsilon^+(\delta^{(i)}, \beta - \delta^{(i)})^{-1} d_i(b) d^{\beta-\delta^{(i)}} \pmod{W}. \tag{3.17}$$

By Lemma 3.4, $d_1, d_2, \dots, d_n$ are A -linear independent derivations. By Lemma 3.7, $Ad^{\beta-\delta^{(i)}} \subset W$ for $i = 1, 2, \dots, n$. This in particular proves i), and that $\oplus_{\alpha \in \mathcal{J} \setminus \{\gamma\}} Ad^\alpha \subset W$ if $|\mathcal{J}| < \infty$. Next we assume that $|\mathcal{J}| < \infty$. Then $|\mathcal{I}| = m < \infty$ and either $\text{char } F = p > 0$ (in this case $\gamma_i = p - 1$ or 1 for $i \in \mathcal{I}$) or $\mathcal{D} = \mathcal{D}_-$ (in this case $\gamma_i = 1$ for $i \in \mathcal{I}$). Then $\mathcal{D}(A)d^\gamma = [\mathcal{D}, Ad^\gamma] \subset W$. Thus $W' = \oplus_{\alpha \in \mathcal{J} \setminus \{\gamma\}} Ad^\alpha \oplus \mathcal{D}(A)d^\gamma \subset W$. To prove $W \subset W'$, let $\alpha, \beta \in \mathcal{J}$ be such that $\alpha + \beta - \gamma \in \mathcal{J} \setminus \{0\}$. It suffices to show the coefficient of d^γ in $[xd^\alpha, yd^\beta]$ to be in $D(A)$. By (2.7), (2.10), the coefficient of d^γ in $[ud^\alpha, vd^\beta]$ for $u, v \in H(A)$ is

$$\begin{aligned}
&(\frac{\alpha}{\gamma-\beta}) \epsilon^+(\alpha + \beta - \gamma, \gamma - \beta)^{-1} \epsilon(d^{\gamma-\beta}, v) \epsilon^+(\gamma - \beta, \beta) u d^{\alpha+\beta-\gamma}(v) \\
&\quad - \epsilon(ud^\alpha, vd^\beta) (\frac{\beta}{\gamma-\alpha}) \epsilon^+(\alpha + \beta - \gamma, \gamma - \alpha)^{-1} \epsilon(d^{\gamma-\alpha}, u) \epsilon^+(\gamma - \alpha, \alpha) v d^{\alpha+\beta-\gamma}(u) \\
&= \epsilon^+(\alpha + \beta - \gamma, \gamma - \beta)^{-1} \epsilon(d^{\gamma-\beta}, v) \epsilon^+(\gamma - \beta, \beta) \left((\frac{\alpha}{\gamma-\beta}) u d^{\alpha+\beta-\gamma}(v) \right. \\
&\quad \left. - (\frac{\beta}{\gamma-\alpha}) \epsilon(\bar{u}, \beta + \alpha - \gamma) d^{\alpha+\beta-\gamma}(u) v \right).
\end{aligned} \tag{3.18}$$

Noting that $\text{char} F = p > 0$ or $\gamma_i = 1$, from the condition $\alpha, \beta, \gamma - \alpha - \beta \in \mathcal{J} \setminus \{0\}$ we see that

$$(\alpha_{\gamma-\beta}) = (\alpha_{\alpha+\beta-\gamma}) = (-1)^{|\alpha+\beta-\gamma|} (= \beta_{\gamma-\alpha}). \quad (3.19)$$

Applying d_i to $u'v'$ for any $u'v' \in H(A)$, we have

$$d_i(u')v' = -\epsilon(u', d_1)u'd_i(v') \in D(A).$$

Using this and (3.19), one can easily deduce that the right-hand side of (3.18) is in $D(A)$. This proves ii). To prove iii), we use Lemma 3.7. By i) and ii), $Z = F_1$ and we have $\dim_{Z(A[D])} A[D] > 4$ except when $A = F_1 \oplus F_1x$ with $x^2 = 0$, and $D = F \frac{\partial}{\partial x}$. This leads to the iii). $\square$

Example 3.10. Let $n \geq 2$ and let $x_1, x_2, \dots, x_n$ be n symbols which have colors in Γ_- . Let A be the free ϵ -commutative associative algebra generated by $x_1, x_2, \dots, x_n$. Clearly A has dimension 2^n . Let D be the space spanned by the derivations $\partial_i = \frac{\partial}{\partial x_i}$, $i = 1, 2, \dots, n$. Then we obtain a simple Lie color algebra $\overline{A[D]}$ of dimension $2^{2n} - 2$. In particular, if we take $\Gamma = \mathbb{Z}/2\mathbb{Z}$, $\epsilon(i, j) = (-1)^{ij}$, $i, j \in \mathbb{Z}/2\mathbb{Z}$, then $A = \Lambda^n(x_1, x_2, \dots, x_n)$ is the exterior algebra and we obtain the simple Hamiltonian Lie superalgebra $\overline{A[D]} = H(2n)$ (see §3.3 in [K]).

References

- [J] D.A. Jordan, On the simplicity of Lie algebras of derivations of commutative algebras, *J. Alg.*, **228** (2000), 580-585.
- [K] V.G. Kac, Lie superalgebras, *Adv. Math.*, **26** (1977), 8-96.
- [O] J.M. Osborn, New simple infinite-dimensional Lie algebras of characteristic 0, *J. Alg.*, 185(1996), no.3, 820-835.
- [P1] D.S. Passman, Simple Lie algebras of Witt type, *J. Alg.*, **206** (1998), 682-692.
- [P2] D.S. Passman, Simple Lie color algebras of Witt type, *J. Alg.*, **208** (1998), 698-721.
- [SZ1] Y. Su, K. Zhao, Simple algebras of Weyl type, *Science in China A*, **44**(2001), 419-426.
- [SZ2] Y. Su, K. Zhao, Isomorphism classes and automorphism groups of algebras of Weyl type, *Science in China A*, **31**(2001), 961-972.
- [Z1] K. Zhao, Simple algebras of Weyl type, II, *Proc. Amer. Math. Soci.*, **130**(2002), 1323-1332.
- [ZS] K. Zhao, Y. Su, Simple Lie color algebras from graded associative algebras, to appear.

Addresses: Department of Mathematics, Shanghai Jiaotong University, Shanghai 200030, China. Email: kfmimi@public1.sta.net.cn.

Academy of Mathematics and system Sciences, Chinese Academy of Sciences, Beijing 100080, China. Email: kzhao@math08.math.ac.cn.

Department of Mathematics, Nanjing University, Nanjing 210093, China. Email: lszhu@pub.sz.jsinfo.net.